BE SMART!

CONCUSSION AWARENESS & MANAGEMENT

WHEN IN DOUBT - SIT THEM OUT!

CONCUSSION: Signs & Symptoms

- Confusion and Disorientation
- Loss of Consciousness
- Headache
- Dizziness
- Nausea and Vomiting
- Loss of Balance

- Double Vision or Fuzzy Vision
- Ringing in the Ears
- Slow or Slurred Speech
- Seeing "Stars"
- Feeling Stunned or Dazed
- Emotional or Personality Changes

CONCUSSION: Management & Rehabilitation

An athlete should never return to play while symptomatic. When a player shows ANY SYMPTOMS or SIGNS of a concussion...

The player should not be allowed to return to play in the current game or practice.

The player should not be left alone, regular monitoring for deterioration is essential.

The player should be medically evaluated.

CONCUSSION: Guidelines for Coaches, Players, Parents & Officials

- Concussion is a Brain Injury
- You do not have to lose consciousness to have a concussion
- Symptoms are often subtle
- Wear properly fitted protective equipment
- The head (helmet/facemask) should never be used to make initial contact with another player
- A concussion may be caused by a direct blow to the head, face, neck, or anywhere else that causes a severe and sudden movement to the head/neck
- Medical Clearance by an appropriate physician is mandatory before return to play

CONCUSSION: Follow these 6 steps before returning to play.

Concussion Management should adhere to the following 6 steps:

- 1. No activity, complete rest. Once asymptomatic, proceed to step two
- 2. Light aerobic exercise such as walking or stationary cycling
- **3.** Sport-specific training. (e.g. running in football, skating in hockey)
- 4. Non-contact training drills
- **5.** Full-contact training after medical clearance
- **6.** Game Play









